

REMARKS

I. Status of the Claims

Claims 1-5 and 9-25 are pending in the application. Claims 1, 3-5, 9-21 and 23-25 stand rejected, variously, under 35 U.S.C. §112, first and second paragraphs, and 35 U.S.C. §102. The specific grounds for rejection, and applicants' response thereto, are set out in detail below.

New claims 36-43 are provided and are supported by the original claims.

II. Rejection Under 35 U.S.C. §112, First Paragraph

Claims 1, 3-5, 9-21 and 23-25 stand rejected under the first paragraph of §112 as lacking enablement. According to the examiner, while the claims are enabled for protecting a mouse from an organophosphate comprising administering to the mouse an expression construct comprising a CMV promoter linked to a PON1 gene, the specification does not provide enablement using promoters generally. Applicants traverse.

The examiner's only evidentiary support for the argument that not all promoters will work derives from Furlong *et al.*, which support is misplaced. Furlong can be distinguished in two ways from the present claims. First, Furlong was in fact seeking to make transgenic animals. That is *not* applicants' intention here, as exemplified by the fact that their examples show a *non-integrative* virus (adenovirus). Thus, to suggest that the promoter selection for the present claims must support long-term stable expression, as Furlong was attempting, simply is incorrect. Second, Furlong's paper was published in 1994, some 7 years prior to the instant application's priority date. And third, Furlong's statements regarding the lack of understanding regarding PON1 regulatory mechanisms and cis-acting elements is similarly irrelevant since

applicants' choice of heterologous promoters avoids such issues, which would be associated with the *native* PON1 promoter.

Thus, it is respectfully submitted that the examiner has not made out a *prima facie* case of lack of enablement for promoters generally. Reconsideration and withdrawal of the rejection is therefore respectfully requested.

III. Rejection Under 35 U.S.C. §112, Second Paragraph

Claims 1-5 and 9-25 stand rejected as allegedly indefinite in failing to recite all necessary steps. It is argued that the step of exposing the cell or subject to an organophosphate is required. Applicants traverse. Because the organophosphate is a toxin, there is no intentional "exposing" – this may or may not happen as a result of the cell or subject entering an environment where the organophosphate exists or appears. Those persons providing the PON1 gene would not be the same persons (if any) introducing the organophosphate. Thus, what the examiner suggests for the claim to be "complete" is, in fact, illogical. That said, applicants have amended the wherein clauses of claims 1 and 21 so that they track more closely with and fully satisfy the claim preamble. Reconsideration and withdrawal of the rejection is therefore respectfully requested.

IV. Rejections Under 35 U.S.C. §102 and §103

Claims 1-5, 91-5 and 17-25 are rejected as anticipated by, and claims 1 and 19 are rendered obvious by (in conjunction with Mackness *et al.*), Hudson *et al.* (U.S. Patent 5,629,193). Applicants traverse.

According to the examiner, Hudson *et al.* teaches the administration of a PON1 gene using a variety of vectors, for the purpose of preventing cell death due to organophosphate toxicity. However, the examiner is incorrect in citing Hudson *et al.* as teaching PON1. Hudson

et al. teaches **PON2**, which explains why they advanced and secured composition of matter claims – indeed, had they been attempting to claim PON1 sequences, they would have failed because PON1 was previously described in the literature (see Hassett *et al.*, “Characterization of cDNA clones encoding rabbit and human serum paraoxonase: the mature protein retains its signal sequence,” *Biochemistry* 30(42):10141-9, Oct 22, 1991; attached). What Hudson *et al.* failed to appreciate at the time of filing, likely because they had obtained **absolutely no data whatsoever**, was that they had cloned an enzyme that operated quite differently than PON1. In any event, Hudson *et al.* cannot anticipate the claims of the present application because the sequence provided in SEQ ID NOS:1 and 2 and in FIG. 1A-B is **not** PON1, but PON2 (see attached sequence information).


Turning to the obviousness rejection, applicants would direct the examiner to statements made in the first office action, mailed on August 23, 2005, where it was stated that as of 2001-2002, gene therapy was “unpredictable.” If gene therapy was unpredictable in 2001-2002, then it was nearly **unattainable** as of 1994, when Hudson *et al.* was filed. In this regard, the complete and utter absence of **any** data in Hudson *et al.* regarding the ability to transfer nucleic acids into living organisms and provide protection is a glaring deficiency that cannot be overlooked. To suggest that the scant disclosure provided by Hudson *et al.* renders obvious applicants’ invention, which is supported by *in vivo* data, contradicts the examiner’s stance on the record regarding gene therapy. Furthermore, reliance on Mackness might provide a correct citation to PON1, but it fails to establish that one could achieve, via gene therapy, protection of subjects from organophosphate toxins. As such, it is submitted that the obviousness rejection cannot stand.

V. Conclusion

In light of the foregoing, applicants submit that all claims are in condition for allowance, and an early notification to that effect is earnestly solicited. Should the examiner have any questions regarding this response, a telephone call to the undersigned is invited.

Please date stamp and return the enclosed postcard as evidence of receipt.

Respectfully submitted,



Steven L. Highlander
Reg. No. 37,642
Attorney for Applicants

FULBRIGHT & JAWORSKI L.L.P.
600 Congress Avenue, Suite 2400
Austin, Texas 78701
(512) 474-5201

Date: February 6, 2007

Characterization of cDNA Clones Encoding Rabbit and Human Serum Paraoxonase: The Mature Protein Retains Its Signal Sequence^{†,‡}

Christopher Hassett,[§] Rebecca J. Richter,^{§,||} Richard Humbert,^{§,||} Christine Chapline,[°] John W. Crabb,[°] Curtis J. Omiecinski,^{§,||} and Clement E. Furlong^{*,||,||}

Department of Environmental Health, School of Public Health and Community Medicine, Department of Genetics, Division of Medical Genetics, School of Medicine, and Dana Program in Ecogenetics, University of Washington, Seattle, Washington 98195, and The W. Alton Jones Cell Science Center, Lake Placid, New York 12946

Received April 5, 1991; Revised Manuscript Received July 1, 1991

ABSTRACT: Serum paraoxonase hydrolyzes the toxic metabolites of a variety of organophosphorus insecticides. High serum paraoxonase levels appear to protect against the neurotoxic effects of organophosphorus substrates of this enzyme [Costa et al. (1990) *Toxicol. Appl. Pharmacol.* 103, 66-76]. The amino acid sequence accounting for 42% of rabbit paraoxonase was determined by (1) gas-phase sequencing of the intact protein and (2) peptide fragments from lysine and arginine digests. From these data, two oligonucleotide probes were synthesized and used to screen a rabbit liver cDNA library. A clone was isolated and sequenced, and contained a 1294-bp insert encoding an open reading frame of 359 amino acids. Northern blot hybridization with RNA isolated from various rabbit tissues indicated that paraoxonase mRNA is synthesized predominantly, if not exclusively, in the liver. Southern blot experiments suggested that rabbit paraoxonase is coded by a single gene and is not a family member of closely related genes. Human paraoxonase clones were isolated from a liver cDNA library by using the rabbit cDNA as a hybridization probe. Inserts from three of the longest clones were sequenced, and one full-length clone contained an open reading frame encoding 355 amino acids, four less than the rabbit paraoxonase protein. Each of the human clones appeared to be polyadenylated at a different site, consistent with the absence of the canonical polyadenylation signal sequence. Of potential significance with respect to the paraoxonase polymorphism, the derived amino acid sequence from one of the partial human cDNA clones differed at two positions from the full-length clone. Amino-terminal sequences derived from purified rabbit and human paraoxonase proteins suggested that the signal sequence is retained, with the exception of the initiator methionine residue [Furlong et al. (1991) *Biochemistry* (preceding paper in this issue)]. Characterization of the rabbit and human paraoxonase cDNA clones confirms that the signal sequences are not processed, except for the N-terminal methionine residue. The rabbit and human cDNA clones demonstrate striking nucleotide and deduced amino acid similarities (greater than 85%), suggesting an important metabolic role and constraints on the evolution of this protein.

Polymorphic genes encoding human biotransformation enzymes which result in variable rates of metabolism of certain drugs and xenobiotics have been identified. Examples of polymorphic enzymes include cytochrome P450 isozymes which hydroxylate the antihypertensive drug debrisoquine and the anticonvulsant mephenytoin [Kalow, 1987], an N-acetyltransferase which metabolizes arylamine and hydrazine compounds [Weber, 1987], the glutathione transferase μ isozyme which conjugates glutathione to electrophilic compounds [Seidegard et al., 1988], and serum cholinesterase which metabolizes the anesthetic succinylcholine [Brown et al., 1981].

Paraoxonase, like serum cholinesterase, demonstrates a substrate-dependent polymorphism in human populations [see

Geldmacher-von Malinckrodt and Diepgen (1988) for review]. Some paraoxonase substrates, such as phenylacetate and chlorpyrifos oxon, are hydrolyzed with the same turnover number by both allelic forms of the enzyme, whereas paraoxon is hydrolyzed slowly by one allelic form and rapidly by the other [LaDu et al., 1986; Furlong et al., 1989; Smolen et al., 1991]. It has been suggested that high serum levels of paraoxonase may be protective against poisoning by organophosphate substrates of this enzyme [Omenn, 1987; LaDu & Eckerson, 1984; Furlong et al., 1988, 1989]. Experiments with animal systems support this hypothesis [Main, 1956; Costa et al., 1990].

One of our aims is to determine the molecular basis for the paraoxonase polymorphism observed in humans. Because rabbits have very high levels of paraoxonase [Costa et al., 1987], we first purified and partially sequenced rabbit paraoxonase [Furlong et al., 1991]. The protein sequence data were used to design oligonucleotide probes which permitted the isolation of a rabbit cDNA. The rabbit clone was subsequently used as a probe to isolate human paraoxonase cDNAs. This report describes these cloning experiments and presents the structural characterization of rabbit and human paraoxonase.

MATERIALS AND METHODS

Protein Purification. Paraoxonase was purified through the DEAE-cellulose fractionation step as described previously [Furlong et al., 1991]. Paraoxonase was further purified by

[†]Supported by NIH Grants GM-15253 (Center Grant), ES-05194 (to C.E.F.), GM-32281, ES-04696, and ES-04978 (to C.J.O.), and EY-06603 and CA-37589 (to J.W.C.) and by a grant from the Charles A. Dana Foundation.

[‡]The nucleotide sequences in this paper have been submitted to the GenBank/EMBL Data Bank under Accession Numbers M63011, M63012, M63013, and M63014.

[§]Address correspondence to this author at the Department of Genetics, SK-50, University of Washington, Seattle, WA 98195.

^{||}Department of Environmental Health, School of Public Health and Community Medicine, University of Washington.

[°]Department of Genetics, University of Washington.

^{||}Division of Medical Genetics, School of Medicine, University of Washington.

[°]Dana Program in Ecogenetics, University of Washington.

[°]The W. Alton Jones Cell Science Center.

high-performance chromatography on a 5- μ m Vydac C₁₈ column.

Protein Digests and Peptide Purification. Paraoxonase was pyridylethylated and succinylated as described by Crabb et al. (1988). Pyridylethylated HPLC-purified paraoxonase was fragmented at lysyl residues with endoproteinase Lys-C (Crabb et al., 1986). Pyridylethylated, succinylated DEAE-purified paraoxonase was cleaved at arginyl residues with trypsin (Crabb et al., 1986). Peptides were purified by narrow-bore reverse-phase HPLC using an Applied Biosystems Model 130 HPLC system.

Protein/Peptide Sequencing. Intact paraoxonase and fractionated peptides were sequenced with an Applied Biosystems gas-phase sequencer (Model 470) and an on-line phenylthiohydantoin amino acid analyzer (Model 120) using the 03RPTH sequencer program and the manufacturer's recommended program and solvents for the PTH analyzer (Crabb et al., 1988). Phenylthiocarbonyl (PTC) amino acid analysis was performed according to West and Crabb (1990) using an Applied Biosystems automatic system (Models 420H/130/920).

Oligonucleotide Synthesis. DNA probes and primers were synthesized with an Applied Biosystems DNA synthesizer using phosphoramidite chemistry.

Library Screening and Subcloning. (A) *Rabbit.* A λ gt11 cDNA library constructed from the pooled livers of male and female New Zealand white rabbits was obtained from Clontech (Palo Alto, CA). The library was screened as described previously (Hassett & Omiecinski, 1987; Hassett et al., 1989) by using the oligonucleotide probes described under Results. The rabbit insert was subcloned into pUC13 with *Escherichia coli* DH5 α as host (BRL, Gaithersburg, MD).

(B) *Human.* A λ gt11 human liver cDNA library derived from an adult female was also obtained from Clontech. This library was screened with the 952-bp *Bst*XI restriction fragment from rabbit paraoxonase cDNA. Inserts were subcloned in pSK(+) Bluescript plasmid vector and used to transform XL1-Blue cells (Stratagene, La Jolla, CA).

DNA Sequence Analysis. (A) *Rabbit.* The insert cDNA was sequenced directly in pUC13 by using the forward and reverse universal plasmid primers and the 17-base paraoxonase-specific primer. Insert DNA was subcloned into the vector in both orientations, relative to the multiple cloning site. Unique *Bam*HI and *Hind*III restriction sites in the insert DNA and in the vector cloning region allowed deletion constructs to be engineered which facilitated sequence analysis of both strands from the universal primers. Each DNA strand was sequenced at least three times.

(B) *Human.* The nucleotide sequence of the human DNA clones was determined in the plasmid by using primers complementary to the T3 and T7 promoters of the vector. Additionally, 11 oligonucleotide primers were synthesized for sequencing on the basis of the derived human and rabbit sequences.

DNA was sequenced by using the dideoxy termination method (Sanger et al., 1977) and Sequenase Version 2.0 (U.S. Biochemicals, Cleveland, OH), as described previously (Hassett & Omiecinski, 1990). Sequence analysis and database searches were performed with either GENEPRO (Riverside Scientific Enterprises, Bainbridge Island, WA) or Intelligenetics (Palo Alto, CA) software and databases, which included GenBank and EMBL DNA databases and the PIR protein database.

Northern Blot Analysis. RNA was isolated (Omiecinski et al., 1985) from the liver, lung, kidney, and testes of two New




FIGURE 1: Reverse-phase HPLC purification of rabbit paraoxonase. Paraoxonase was purified through the DEAE-Trisacryl M step (87 μ g) and fractionated by reverse-phase HPLC on a 5- μ m Vydac C₁₈ column. Solvent A was 0.1% trifluoroacetic acid in H₂O, and solvent B was 84% acetonitrile containing 0.09% trifluoroacetic acid.

Zealand White rabbits. Twenty micrograms of total RNA from each organ was size-fractionated in a 6% formaldehyde/1.15% agarose gel and transferred to a GeneScreen Plus nylon membrane as per the manufacturer's directions (Du Pont/NEN, Boston, MA). A 438-bp *Bam*HI fragment isolated from the 3' region of the rabbit paraoxonase cDNA was radiolabeled (Hassett & Omiecinski, 1990) and used as a hybridization probe. The membrane was washed at 45 $^{\circ}$ C in 1.0 \times SSC/0.1% SDS (1 \times SSC: 1.5 M NaCl, 0.15 M sodium citrate) and exposed overnight to X-ray film in the presence of two intensifying screens. The size of the *in vivo* RNA transcript was estimated by using an RNA ladder standard (BRL, Gaithersburg, MD).

Southern Blot Analysis. Peripheral white blood cell DNA was extracted and isolated from 5 mL of whole blood withdrawn from a single rabbit and processed essentially as described (Blin & Stafford, 1976). Twenty micrograms of DNA was digested with *Eco*RI, *Bam*HI, *Hind*III, *Pst*I, or *Xho*I, size-fractionated on a 0.85% agarose gel, and transferred to a nylon membrane as described previously (Hassett et al., 1989). The Southern blot was incubated with a radiolabeled 419-bp fragment isolated from the rabbit paraoxonase cDNA (*Eco*RI/*Bam*HI fragment). The blot was washed in a final solution of 0.1 \times SSC/0.1% SDS at 50 $^{\circ}$ C and exposed to X-ray film for 6 days in the presence of two intensifying screens. Drigert III (Pharmacia, Piscataway, NJ) was employed as a molecular size standard.

RESULTS

Purification of Rabbit Paraoxonase. Rabbit paraoxonase, purified through the DEAE-Trisacryl M step as described in the preceding paper, was further purified by high-performance reverse-phase liquid chromatography (Figure 1). Peak 15 contained only homogeneous paraoxonase while peaks 22 and 23 contained both paraoxonase and apolipoprotein A1 (de-

Table I: Rabbit Paraoxonase Protein/Peptide Sequence^a

protein/peptide	sequence	position
amino terminus	AKLTALTLLGLGLALFDGQKS-FQT	2-26
lysyl peptides	S-FQTRFNVHREVPVLPN-NL	22-44
	<u>L</u> PSVNDIVAVGPEHYFA	163-180
	IHYVEK	245-250
	SIDFNTLVDSISVDPV	261-276
	NPPASEVLRIQDIL	298-311
	ALY-ELSQAN	350-359
arginyl peptides	FNVHR	28-32
	VVAEGDFDANGINISPDGKYVYIAELLAHKI-VY	215-248
	IFYYDPKNPPASEVLR	291-306
	IQDILSKPEKV-VAYAE	307-323

^aCycles where no residue was assigned are shown as dashes. Tentative assignments are underlined. The single assignment which differed from the deduced sequence shown in Figure 3 is italicized. Sequences that were used for probe design are shown in bold type.




FIGURE 2: HPLC purification of lysine peptides from paraoxonase. Peptides resulting from the cleavage of pyridylethylated paraoxonase (~78 μ g) with endoprotease Lys-C were purified by narrow-bore reverse-phase HPLC with a linear gradient of buffer A (0.1% TFA) to 50% buffer B (5% acetonitrile, 0.005% TFA) run over 40 min.

terminated by sequence analysis).

Peptide Generation. RP-HPLC-purified pyridylethylated paraoxonase was digested with endoproteinase Lys-C, and the resulting peptides were purified by narrow-bore RP-HPLC (Figure 2) as described under Materials and Methods. In addition, paraoxonase (200 μ g) purified by DEAE-Trisacryl M chromatography (preceding paper) was succinylated and then subjected to arginyl-specific cleavage with trypsin, and the resulting peptides were purified by narrow-bore RP-HPLC. Since the intact paraoxonase had not been RP-HPLC-purified prior to digestion, peptides from apolipoprotein A1 were also identified by sequence analysis (data not shown).

Gas-Phase Protein Sequencing. Unequivocal amino acid sequencing data were obtained from rabbit paraoxonase, four arginyl peptides, and six lysyl peptides (Table I). Residues 16-20 of the amino terminus of rabbit paraoxonase (Phe-Asp-Gly-Gln-Lys) allowed the design of the 15-base oligo-nucleotide 5'-TTY GAY GGN CAR AAR-3' with 64-fold redundancy (Table I, Figure 3). A 17-base oligomer (5'-GGG TCR TAR TAR AAD AT-3') with 48-fold redundancy was designed as a complement to the nucleotide sequence encoding residues 1-6 of the arginine peptide IFYYDP (Table I, Figure 3). The wobble position of the proline codon was not used in the design of this oligomer.

Isolation and Sequence of Rabbit Paraoxonase cDNA. Approximately 400,000 plaques were screened from the rabbit cDNA library with the 15-base probe, yielding 35 potentially positive autoradiographic signals. Twenty-four of these plaques were rescreened with the 17-base probe, and a plaque which

hybridized to this oligomer was plated at a low density. This cDNA clone was screened a final time with the 15-base probe and once again showed positive hybridization. Phage DNA was purified, digested with *Eco*RI, and subcloned into pUC13.

DNA sequence analysis of the rabbit paraoxonase cDNA (RabPON, GenBank Accession Number M63011) identified an insert of 1294 bp, containing the entire protein coding sequence (Figure 3). Fifty-one nucleotides precede the methionine initiation codon, ATG, which begins an open reading frame coding for 359 amino acids. An amber stop codon, TAG, is followed by an additional 163 nucleotides of 3' non-coding sequence. The ATG at position 1 is the likely start position since there is a stop codon beginning 15 nucleotides upstream from this ATG. No poly(A) signal or sequence was identified in this clone. The fragment containing this information was presumably deleted during library construction since multiple efforts to isolate this region from the original λ phage were unsuccessful.

Comparison between the Derived and Determined Amino Acid Sequences. The deduced amino acid sequence is shown in Figure 3. The sequence verified by gas-phase amino acid sequencing is presented in Table I. The verified protein sequence totaled 151 residues or about 42% of the rabbit paraoxonase protein sequence deduced from the cDNA clone. The one difference observed (i.e., Ile for Leu at position 164) may simply reflect a variant in the rabbit population.

Analysis of Paraoxonase mRNA Expression in Rabbit Tissues. Northern blot analysis performed with RNA isolated from four rabbit organs revealed the presence of paraoxonase-specific RNA in liver only. RNA isolated from lung, kidney, or testes did not hybridize to the paraoxonase cDNA probe (Figure 4). On the basis of this Northern blot and linear regression analysis, the molecular size estimate for the *in vivo* liver mRNA transcript was approximately 1400 bases. Pretreatment of animals with phenobarbital 16 h prior to sacrifice did not influence steady-state mRNA levels of liver paraoxonase (data not shown).

Evaluation of Rabbit Paraoxonase Gene Complexity. Southern-blotted rabbit genomic DNA was digested with five restriction endonucleases prior to electrophoresis and probed with the 400-bp *Eco*RI/*Bam*HI fragment of the rabbit paraoxonase cDNA. In each restriction digest lane, only one hybridization band was observed (Figure 5). These data suggest that rabbit paraoxonase protein is probably encoded by a single gene, and not a member of a family of closely related genes.

Isolation and Sequence of Human Paraoxonase cDNAs. Approximately 300,000 plaques from the human cDNA library were screened with a 952-bp *Bst*XI radiolabeled fragment from the rabbit paraoxonase cDNA. From this library screen, 41 plaques were identified, and the three longest clones were

¹ As recommended by the Nomenclature Committee of the International Union of Biochemistry, nucleotides are abbreviated as follows: R = purine; Y = pyrimidine; N = A, T, G, or C; D = G, A, or T.

RabPON															CGG CCC			-46
AGC	CCG	TGG	TGC	TCG	CGC	CGG	TCC	AGC	CTT	TAG	TCT	GCC	CTG	ACC	-1			
HuPONI												CCC	CCG	ACC	-1			
ATG	GCT	AAA	CTG	ACA	GCG	CTC	ACG	CTC	TTG	GGG	CTG	GGA	TTG	GCA	45			
Met	Ala	Lys	Leu	Thr	Ala	Leu	Thr	Leu	Leu	Gly	Leu	Gly	Leu	Ala				
Met	Ala	Lys	Leu	Ile	Ala	Leu	Thr	Leu	Leu	Gly	Met	Gly	Leu	Ala	15			
ATG	GCG	AAG	CTG	ATT	GCG	CTC	ACC	CTC	TTG	GGG	ATG	GGA	CTG	GCA				
CTC	TTC	GAT	GGA	CAG	AAG	TCT	TCT	TTC	CAA	ACA	CGA	TTT	AAT	GTT	90			
Leu	Phe	Asp	Gly	Gln	Lys	Ser	Ser	Phe	Gln	Thr	Arg	Phe	Asn	Val				
Leu	Phe	Arg	Asn	His	Gln	Ser	Ser	Tyr	Gln	Thr	Arg	Leu	Asn	Ala	30			
CTC	TTC	AGG	AAC	CAC	CAG	TCT	TCT	TAC	CAA	ACA	CGA	CTT	AAT	GCT				
CAC	CGT	GAA	GTA	ACT	CCA	GTG	GAA	CTT	CCT	AAC	TGT	AAT	TTA	GTT	135			
His	Arg	Glu	Val	Thr	Pro	Val	Glu	Leu	Pro	Asn	Cys	Asn	Leu	Val				
Leu	Arg	Glu	Val	Gln	Pro	Val	Glu	Leu	Pro	Asn	Cys	Asn	Leu	Val	45			
CTC	CGA	GAG	GTA	CAA	CCC	GTA	GAA	CTT	CCT	AAC	TGT	AAT	TTA	GTT				
AAA	GGG	ATT	GAC	AAT	GGT	TCT	GAA	GAC	TTG	GAA	ATA	CTG	CCC	AAT	180			
Lys	Gly	Ile	Asp	Asn	Gly	Ser	Glu	Asp	Leu	Glu	Ile	Leu	Pro	Asn				
Lys	Gly	Ile	Glu	Thr	Gly	Ser	Glu	Asp	Met	Glu	Ile	Leu	Pro	Asn				
AAA	GGA	ATC	GAA	ACT	GGC	TCT	GAA	GAC	ATG	GAG	ATA	CTG	CCT	AAT	60			
GGA	CTG	GCT	TTC	ATC	AGC	GCC	GGA	TTA	AAA	TAT	CCT	GGA	ATA	ATG	225			
Gly	Leu	Ala	Phe	Ile	Ser	Ala	Gly	Leu	Lys	Tyr	Pro	Gly	Ile	Met				
Gly	Leu	Ala	Phe	Ile	Ser	Ser	Gly	Leu	Lys	Tyr	Pro	Gly	Ile	Lys	75			
GGA	CTG	GCT	TTC	ATT	AGC	TCT	GGA	TTA	AAG	TAT	CCT	GGA	ATA	AAG				
AGC	TTT	GAC	CCT	GAT	AAG	CCT	GGA	AAG	ATA	CTT	CTA	ATG	GAC	CTG	270			
Ser	Phe	Asp	Pro	Asp	Lys	Pro	Gly	Lys	Ile	Leu	Leu	Met	Asp	Leu				
Ser	Phe	Asn	Pro	Asn	Ser	Pro	Gly	Lys	Ile	Leu	Leu	Met	Asp	Leu	90			
AGC	TTT	GAC	CCC	AAC	AGT	CCT	GGA	AAA	ATA	CTT	CTG	ATG	GAC	CTG				
AAT	GAG	AAA	GAC	CCA	GTA	GTA	TTG	GAA	CTG	AGC	ATT	ACT	GGA	AGT	315			
Asn	Glu	Lys	Asp	Pro	Val	Val	Leu	Glu	Leu	Ser	Ile	Thr	Gly	Ser				
Asn	Glu	Glu	Asp	Pro	Thr	Val	Leu	Glu	Leu	Gly	Ile	Thr	Gly	Ser	105			
AAT	GAA	CAA	GAT	CCA	ACA	GTG	TTG	GAA	TTG	GGG	ATC	ACT	GGA	AGT				
ACA	TTT	GAT	TTA	TCT	TCA	TTT	AAC	CCT	CAT	GGG	ATT	AGC	ACA	TTC	360			
Thr	Phe	Asp	Leu	Ser	Ser	Phe	Asn	Pro	His	Gly	Ile	Ser	Thr	Phe				
Lys	Phe	Asp	Val	Ser	Ser	Phe	Asn	Pro	His	Gly	Ile	Ser	Thr	Phe	120			
AAA	TTT	GAT	GTA	TCT	TCA	TTT	AAC	CCT	CAT	GGG	ATT	AGC	ACA	TTC				
ACA	GAT	GAA	GAT	AAT	ATC	GTC	TAC	CTG	ATG	GTG	GTG	AAC	CAT	CCA	405			
Thr	Asp	Glu	Asp	Asn	Ile	Val	Tyr	Leu	Met	Val	Val	Asn	His	Pro				
Thr	Asp	Glu	Asp	Asn	Ala	Met	Tyr	Leu	Leu	Val	Val	Asn	His	Pro	135			
ACA	GAT	GAA	GAT	AAT	CCC	ATG	TAC	CTC	CTG	GTG	GTG	AAC	CAT	CCA				
GAT	TCC	AAG	TCC	ACA	GTG	GAG	TTG	TTT	AAA	TTC	CAA	GAA	AAA	GAA	450			
Asp	Ser	Lys	Ser	Thr	Val	Glu	Leu	Phe	Lys	Phe	Gln	Glu	Lys	Glu				
Asp	Ala	Lys	Ser	Thr	Val	Glu	Leu	Phe	Lys	Phe	Gln	Glu	Glu	Glu	150			
GAT	CCC	AAG	TCC	ACA	GTG	GAG	TTG	TTT	AAA	TTT	CAA	GAA	CAA	GAA				
AAA	TCA	CTT	TTG	CAT	CTG	AAA	ACC	ATC	AGA	CAC	AAG	CTT	CTG	CCT	495			
Lys	Ser	Leu	Leu	His	Leu	Lys	Thr	Ile	Arg	His	Lys	Leu	Leu	Pro				
Lys	Ser	Leu	Leu	His	Leu	Lys	Thr	Ile	Arg	His	Lys	Leu	Leu	Pro	165			
AAA	TCG	CTT	TTG	CAT	CTA	AAA	ACC	ATC	AGA	CAT	AAA	CTT	CTG	CCT				
AGT	GTG	AAT	GAC	ATT	GTC	GCT	GTG	GGA	CCT	GAA	CAC	TTT	TAT	GCT	540			
Ser	Val	Asn	Asp	Ile	Val	Ala	Val	Gly	Pro	Glu	His	Phe	Tyr	Ala				
Asn	Leu	Asn	Asp	Ile	Val	Ala	Val	Gly	Pro	Glu	His	Phe	Tyr	Gly	180			
AAA	TTG	AAT	GAT	ATT	GTT	GCT	GTG	GGA	CCT	GAG	CAC	TTT	TAT	GGC				
ACC	AAT	GAT	CAC	TAT	TTT	ATT	GAC	CCT	TAC	TTA	AAA	TCC	TGG	GAA	585			
Thr	Asn	Asp	His	Tyr	Phe	Ile	Asp	Pro	Tyr	Leu	Lys	Ser	Trp	Glu				
Thr	Asn	Asp	His	Tyr	Phe	Leu	Asp	Pro	Tyr	Leu	Gln	Ser	Trp	Glu	195			
ACA	AAT	GAT	CAC	TAT	TTT	CTT	GAC	CCC	TAC	TTA	CAA	TCC	TGG	GAG				
ATG	CAT	TTG	GGA	TTA	GCG	TGG	TCA	TTT	GTT	ACT	TAT	TAT	AGT	CCC	630			
Met	His	Leu	Gly	Leu	Ala	Trp	Ser	Phe	Val	Thr	Tyr	Tyr	Ser	Pro				
Met	Tyr	Leu	Gly	Leu	Ala	Trp	Ser	Tyr	Val	Val	Tyr	Tyr	Ser	Pro	210			
ATG	TAT	TTG	GGT	TTA	GCG	TGG	TCG	TAT	GTT	GTC	TAC	TAT	AGT	CCA				
AAT	GAT	GTT	CGA	GTA	GTG	GCA	GAA	GGA	TTT	GAT	TTT	GCT	AAC	GGA	675			
Asn	Asp	Val	Arg	Val	Val	Ala	Glu	Gly	Phe	Asp	Phe	Ala	Asn	Gly				
Ser	Glu	Val	Arg	Val	Val	Ala	Glu	Gly	Phe	Asp	Phe	Ala	Asn	Gly	225			
AGT	GAA	GTT	CGA	GTG	GTG	GCA	GAA	GGA	TTT	GAT	TTT	GCT	AAT	GGA				
ATC	AAC	ATC	TCA	CCA	GAC	GGC	AAG	TAT	GTC	TAT	ATA	GCT	GAA	CTG	720			
Ile	Asn	Ile	Ser	Pro	Asp	Gly	Lys	Tyr	Val	Tyr	Ile	Ala	Glu	Leu				
Ile	Asn	Ile	Ser	Pro	Asp	Gly	Lys	Tyr	Val	Tyr	Ile	Ala	Glu	Leu	240			
ATC	AAC	ATT	TCA	CCC	GAT	GGC	AAG	TAT	GTC	TAT	ATA	GCT	GAG	TTG				

CTG GCT CAT AAG ATC CAT GTG TAT GAA AAG CAC GCT AAT TGG ACT	765
Leu Ala His Lys Ile His Val Tyr Glu Lys His Ala Asn Trp Thr	
Leu Ala His Lys Ile His Val Tyr Glu Lys His Ala Asn Trp Thr	255
CTG GCT CAT AAG ATT CAT GTG TAT GAA AAG CAT GCT AAT TGG ACT	
TTA ACT CCA TTG AAG TCC CTC GAC TTT AAC ACT CTT GTG GAC AAC	810
Leu Thr Pro Leu Lys Ser Leu Asp Phe Asn Thr Leu Val Asp Asn	
Leu Thr Pro Leu Lys Ser Leu Asp Phe Asn Thr Leu Val Asp Asn	270
TTA ACT CCA TTG AAG TCC CTT GAC TTT AAT ACC CTC GTG GAT AAC	
ATA TCC GTG GAT CCT GTG ACA GGG GAC CTT TGG GTT GGT TGT CAT	855
Ile Ser Val Asp Pro Val Thr Gly Asp Leu Trp Val Gly Cys His	
Ile Ser Val Asp Pro Glu Thr Gly Asp Leu Trp Val Gly Cys His	285
ATA TCT GTG GAT CCT GAG ACA GGA GAC CTT TGG GTT GGA TGC CAT	
CCC AAT GGC ATG GGA ATC TTC TAC TAT GAC CCA AAG AAT CCT CCT	900
Pro Asn Gly Met Arg Ile Phe Tyr Tyr Asp Pro Lys Asn Pro Pro	
Pro Asn Gly Met Lys Ile Phe Phe Tyr Asp Ser Glu Asn Pro Pro	300
CCC AAT GGC ATG AAA ATC TTC TTC TAT GAC TCA GAG AAT CCT CCT	
GCA TCA GAG GTG CTT CGA ATC CAG GAC ATT TTA TCC AAA GAG CCC	945
Ala Ser Glu Val Leu Arg Ile Gln Asp Ile Leu Ser Lys Glu Pro	
Ala Ser Glu Val Leu Arg Ile Gln Asn Ile Leu Thr Glu Glu Pro	315
GCA TCA GAG GTG CTT CGA ATC CAG AAC ATT CTA ACA GAA GAA CCT	
AAA GTG ACA GTG GCT TAT GCA GAA AAT GGC ACT GTG TTA CAG GGC	990
Lys Val Thr Val Ala Tyr Ala Glu Asn Gly Thr Val Leu Gln Gly	
Lys Val Thr Gln Val Tyr Ala Glu Asn Gly Thr Val Leu Gln Gly	330
AAA GTG ACA CAG GTT TAT GCA GAA AAT GGC ACA GTG TTG CAA GGC	
AGC ACG GTG GCC GCT GTG TAC AAA GGG AAA ATG CTG GTT GGC ACC	1035
Ser Thr Val Ala Ala Val Tyr Lys Gly Lys Met Leu Val Gly Thr	
Ser Thr Val Ala Ser Val Tyr Lys Gly Lys Leu Leu Ile Gly Thr	345
AGT ACA GTT GGC TCT GTG TAC AAA GGG AAA CTG CTG ATT GGC ACA	
GTG TTC CAC AAA GCT CTC TAC TGT GAG CTC TCA CAG GCC AAT TAG	1080
Val Phe His Lys Ala Leu Tyr Cys Glu Leu Ser Gln Ala Asn ****	359
Val Phe His Lys Ala Leu Tyr Cys Glu Leu ***	355
GTG TTT CAC AAA GCT CTT TAC TGT GAG CTC TAA CAG ACC GAT TTG	
CAC CCG TGC CCG GGA CAC TGG CAC CCA GGA TTT CAA CTG CTT GCC	1125
CAC CCA TGC CAT AAG ACA TGA GGC CAT TAT TTC AAC CGC TTG CCA	
GGC CAC ATT CTT GGG GCC ACA GTG CCC TCG CGC GGA TGA TGG ACA	1170
TAT TCC GAG GAC CCA GTG TTC TTA GCT GAA CAA TGA ATG CTG ACC	
ACC CTA AAT TTG ACA TCA ACT GCA TCG CAG CCT AGA GTG GAT ATG	1215
CTA AAT GTG GAC ATC ATG AAG CAT CAA AGC ACT GTT TAA CTG GGA	
AAG AGT AGG GCT TTT TGA GCG TGA ATT C	1243
GTG ATA TGA TGT GTA GGG CTT TTT TTT GAG AAT ACA CTA TCA AAT	1260
CAG TCT TGG AAT ACT TGA AAA CCT CAT TTA CCA TAA AAA TCC TTC	1305
CTA CTA AAA TGG ATA AAT CAG TTA AAA AAA AA	1337

FIGURE 3: Nucleotide and deduced amino acid sequences of RabPON and HuPON1 cDNAs. The rabbit sequences are presented in normal font in the upper lines; the human sequences are italicized in the lower lines. Alignment begins at the initiation codon ATG, which is arbitrarily designated position 1. Nucleotides preceding this codon are assigned negative numbers. The regions used for oligomer construction in the rabbit sequence are identified by an overline. Amino acid differences between RabPON and HuPON1 are boxed. Potential N-glycosylation sites are shown in bold type.

characterized by DNA sequencing (GenBank Accession Numbers: HuPON1, M63012; HuPON2, M63013; HuPON3, M63014). The DNA sequence of these clones indicated that only HuPON1 was full length (Figure 3). This 1337-bp cDNA, including a 9-base poly(A) tail at the 3' end, contained an open reading frame of 1065 bases that predicted a 355 amino acid protein. Clones HuPON2 and HuPON3 have 5' termini starting 62 and 96 nucleotides downstream from the 5' end of clone HuPON1, respectively. The nucleotide sequences of clones HuPON1 and HuPON2 predict a methionine at position 55 and glutamine at position 192, while clone HuPON3 predicts a protein with a leucine (TTG) at position 55 and an arginine (CGA) at position 192. The former substitution results in the loss of a restriction site (NlaIII) in HuPON3, while the latter substitution creates *AlwI* and *Sau3A* sites in HuPON3.

Comparison of Rabbit and Human Paraoxonase Sequences. Alignment of rabbit and human cDNA coding regions revealed an 86% identity (Figure 3). The protein sequences deduced

from these clones indicated an 85% identity, which increases to 88.7% when conservative amino acid substitutions are considered (Figure 3). The deduced rabbit amino acid sequence contains five potential N-glycosylation sites, whereas the human sequence predicts four possible N-glycosylation sites.

DISCUSSION

The most difficult step in isolating a cDNA clone for human serum paraoxonase has been obtaining sufficient pure enzyme from which to obtain a protein sequence that in turn could be used to design oligomer probes for library screening. We overcame this problem by purifying and partially sequencing paraoxonase from rabbits, which have much higher levels of paraoxonase than humans and for which an activity stain was developed (preceding paper). These sequence data were used to design oligonucleotide probes which enabled the isolation of a rabbit paraoxonase cDNA. The rabbit clone was used to isolate corresponding human liver cDNA clones.




FIGURE 4: Northern blot analysis of rabbit RNA. Total RNA was isolated from the liver, lung, kidney, and testes of untreated rabbits and size-separated in an agarose/formaldehyde gel. Following transfer to a nylon membrane, the blot was probed with the RabPON cDNA. Molecular size standards are shown in the left margin. A single hybridization band is observed only in liver, suggesting an *in vivo* transcript of approximately 1.4 kb.




FIGURE 5: Southern blot analysis of rabbit genomic DNA. Twenty micrograms of DNA was digested under excess conditions with each of the indicated restriction enzymes, size-separated in an agarose gel, and transferred to a nylon membrane which was hybridized with a radioactive RabPON cDNA fragment. Molecular size markers are shown in the left margin.

The full-length rabbit and human clones demonstrate extensive conservation in nucleotide and deduced amino acid

sequences despite the evolutionary distance separating these species. Although the predicted length of the two proteins differs by four amino acids, no gaps were required for the alignment of these sequences. A region of absolute conservation between the two sequences is observed from amino acids 213 to 275. Furthermore, within these 63 residues, three of the four predicted N-glycosylation sites common to the two proteins occur. Although the rabbit and human paraoxonase clones demonstrate significant conservation, other genes or proteins related to paraoxonase were not identified in database searches, despite the fact that many sequences have been described for proteins which perform similar catalytic functions (e.g., esterase activity). On the basis of the unreported partial peptide sequence from human paraoxonase, Gan et al. (1991) also did not identify closely related sequences in database searches.

Comparison of the deduced protein sequences from RabPON and HuPON1 cDNAs to the amino-terminal sequences determined by gas-phase sequencing of the intact proteins reveals a unique feature of paraoxonase. Both the rabbit and human enzymes retain their signal sequences, with only the amino-terminal methionine residues cleaved. Database searches indicate that the N-terminal sequences of rabbit and human paraoxonase show similarity to other protein secretion signal sequences (Figure 6). Conservation of specific amino acids is apparent, but particularly interesting are the conserved three amino acid residues LAL. An inspection of other published signal sequences (Watson, 1984) indicates that many of these also contain this sequence in the hydrophobic core region.

We are unaware of other examples where typical nonmutant signal sequences are retained in mature, secreted proteins. Cleavage of signal sequences appears to follow certain rules (von Heijne, 1983). Both the human and rabbit sequences possess a positively charged amino terminus commonly found in signal sequences, as well as a 9-residue hydrophobic core starting at position 9. In eukaryotes, cleavage typically occurs 5–6 residues from the C-terminal boundary of the hydrophobic core, which would predict a cleavage site for the paraoxonase proteins following residue 22 or 23. Furthermore, accurate processing is thought to exclude certain residues at the –1 and –3 positions, relative to the cleavage site (von Heijne, 1986). On the basis of the cleavage site positions predicted above, the –3 position would be occupied in the paraoxonase protein by Gln₂₀ or Lys₂₁ (rabbit) or by His₂₀ or Glu₂₁ (human). These are “forbidden” residues in the –3 position and may explain why the signal sequence of paraoxonase is retained.

The function of the retained signal sequence is unknown. The “hydrophobic head” of paraoxonase may be important for interaction with the high-density lipoprotein particle with which it is intimately associated. Detergents are required to dissociate paraoxonase from apolipoprotein A1 (Furlong et al., preceding paper; Gan et al., 1991). Hydrophobicity analyses (Figure 7) clearly show the hydrophobic amino termini of rabbit and human paraoxonases, as well as considerable hydrophobic character in the remainder of the proteins.

Comparison of the full-length HuPON1 cDNA with the two human partial clones reveals two interesting features. First, two nucleotide substitutions result in amino acid differences between clones HuPON1 and HuPON2 vs HuPON3. It is not known if either of these substitutions accounts for the differences observed between high- and low-activity paraoxonase allelic forms. In this regard, it is of interest to compare the two amino acid substitutions predicted from clone HuPON3 with the orthologous positions predicted from the

MAKLTALTLLGLGLALFDGOKSSSFQTR Rabbit Paraoxonase
 MAKLTALTLLGLGLALFRNHOSSYQTR Human Paraoxonase
 MQMSPALTCVLGLLFLVFGGSAVHHF Plasminogen Activator inhibitor-1 precursor, Human¹
 MAPRTLLLLSGALALTQWAGSHSMR HLA alpha chain precursor, clone pHLA, Human²
 MAPRTLLLLSGALALTQWAGSHSMR HLA alpha chain precursor, cw3, Human³
 MAPRTLLLLLAGAALTQWAGSHSMR HLA alpha chain precursor histocompatibility antigen, Rabbit⁴
 MAPRTLLLLLAGAALTQYRAGPHSLR H-2 k-d alpha chain precursor class I antigen⁵
 MAKLLALSLSFCFLLLGGCFALREQPQ Legumin A precursor Garden pea⁶
 MGKSKHSCCSLLLLFAGLASGHQVL α amylase 2-precursor Barley⁷
 MAAATTTSPRLILSRQAASSLQCR Fructose 1, 6 biphosphatase precursor wheat⁸
 TAPRTVLLLLSAAALALTQWAGSHSMR CHLA-81 α chain precursor histocompatibility antigen chimpanzee⁹
 ASRLTLLLLSAAALALTQWAGSHSMR Complement C1 inhibitor precursor - Human¹⁰
 ATKTFALLLLSFLAVGLGEGKKEGHFS Gastric inhibitor polypeptide precursor Human¹¹
 MRMLLLSLALGAAYVVAIPTEIPTS Interleukin 5 Human¹²
 MQMSPALTCVLGLLFLVFGGSAVHHF Plasminogen activator inhibitor-1 precursor Human¹³
 MQKLLCSRLVLAALILVLESSVQGY Secretory granule proteoglycan core protein precursor-Human¹⁴
 MAGPRLILLLPTLLALARGFLCALAAQ T cell surface glycoprotein CD7 precursor Human¹⁵
 EAPIVLLLLWLALAPTGSASSEAPP Rabphorin I precursor Rat¹⁶

FIGURE 6: Comparison of the amino-terminal signal sequence regions of rabbit and human paraoxonases with similar signal sequences found in searching the DNA/protein databases. Numbers appearing to the left of the sequences indicate the residue position. Other sequences being at the first residue. Footnotes: (1) Pannekoek et al., 1986; Ginsburg et al., 1986; (2) Malissen et al., 1982; (3) Soderoy et al., 1984; (4) Tykocinski et al., 1984; (5) Kvist et al., 1983; Lalanne et al., 1983; (6) Lyett et al., 1984; (7) Knox et al., 1987; (8) Raines et al., 1988; (9) Meyer et al., 1988; (10) Bock et al., 1986; (11) Takeda et al., 1987; (12) Azuma et al., 1986; (13) Strandberg et al., 1988; (14) Stevens et al., 1988; (15) Aruffo & Seed, 1987; (16) Harnik-Or et al., 1987.




FIGURE 7: Hydrophobicity profiles of rabbit and human paraoxonase. The analysis was performed with GENEPRO software according to the Hopp and Woods (1981) algorithm with a window setting of 12.

rabbit cDNA. Amino acid 55 is a Leu in both sequences, whereas residue 192 is conservatively substituted (Lys in RabPON, Arg in HuPON3). It is tempting to speculate that since rabbits have high-activity paraoxonase, HuPON3 (which shares two similar and potentially important amino acids) might represent the high-activity allele genotype in the human. A gene frequency for the low-activity paraoxonase allele of 0.69 would predict a 43% probability that a given person would

be a heterozygote (Furlong et al., 1989). Therefore, it is not unlikely that the individual from whom the human liver library was constructed was heterozygous for the paraoxonase allele, expressing mRNA for both high- and low-activity forms of the enzyme.

The importance of these amino acid changes could be examined by different approaches. Expressing the human cDNAs in vitro, or site-directed mutagenesis targeting the

HuPON1:	TAACAGACCGGATTTCACCCATGCCATAGAACTGAGGCCATTATTTCAACCGCTTGCCA	60
HuPON2:	TAACAGACCGGATTTCACCCATGCCATAGAACTGAGGCCATTATTTCAACCGCTTGCCA	
HuPON3:	TAACAGACCGGATTTCACCCATGCCATAGAACTGAGGCCATTATTTCAACCGCTTGCCA	

HuPON1:	TATTCGGAGGACCCAGTGTCTTAGCTGAACAATGAATGCTGACCTAAATGTGGACATC	120
HuPON2:	TATTCGGAGGACCCAGTGTCTTAGCTGAACAATGAATGCTGACCTAAATGTGGACATC	
HuPON3:	TATTCGGAGGACCCAGTGTCTTAGCTGAACAATGAATGCTGACCTAAATGTGGACATC	

HuPON1:	ATGAAGCATCAAGACACTGTTTAACTGGGAGTGATATGATGTGTAGGGCTTTTTTTTGG	180
HuPON2:	ATGAAGCATCAAGACACTGTTTAACTGGGAGTGATATGATGTGTAGGGCTTTTTTTTGG	
HuPON3:	ATGAAGCATCAAGACACTGTTTAACTGGGAGTGATATGATGTGTAGGGCTTTTTTTTGG	

HuPON1:	AATACACTACAAATCAGTCTTGAATACCTTGAACCTCATTACATAAAATCCTTC	240
HuPON2:	AATACACTACAAATCAGTCTTGAACCTCATTACATAAAATCCTTC	218
HuPON3:	AATACACTACAAATCAGTCTTGAATACCTTGAACCTCATTACATAAAATCCTTC	

HuPON1:	TCACATAAATGGATAAATCAGTTAAAAA	272
HuPON3:	TCACATAAATGGATAAATCAGTTATGCTCAATATGTCAGATATTAATAACAGTGTGTGACC	300

HuPON3:	CCAAAGTACTTACCTTAAACATGTGTGCTGAAAGCACATGTGTATCGCTGCCTT	360
HuPON3:	GGCATGTCTTGTTCAGAGACACAGGGAGCAGGGTTAGCTCAGCTGTCTTTAGAATCC	420
HuPON3:	AGTACTCACCAGGACTCCAGTTTACAGGCCAGAAACATATGCATTATGAATTCCTCC	480
HuPON3:	CTTACTCCATGACATAGTAAGTCTGACTATGCGAGTCAGCTTACTTACTCCATTTC	540
HuPON3:	CCTTCGATATATGACTTTTTCTCAGTAAATATTAACTGAACTATTCAAAAA	600
HuPON3:	AAAAA	609

FIGURE 8: Comparison of the 3' noncoding portions of the human paraoxonase cDNAs. HuPON1, HuPON2, and HuPON3 are aligned and numbered beginning with the terminating codon TAA, shown in bold type. Identical nucleotide residues in all sequences are indicated with an asterisk below the aligned residue. Potential polyadenylation signal sequences are underlined.

nucleotides encoding these amino acids, could reveal a concordant relationship with substrate-dependent metabolism and the human polymorphism. A more general approach for the identification of genetic alterations relevant to the paraoxonase polymorphism would be to sequence genomic DNA isolated from individuals characterized for high and low activity and to search for structural differences common to each group. Restriction site differences observed between the human sequences should also be useful in this regard.

A second observation in comparing the human clones is the different lengths of the 3' untranslated regions, shown in Figure 8. The sequences are consistent with the existence of mRNAs which are polyadenylated at different sites. The canonical polyadenylation signal (AATAAA) is not found in any of these clones, although potential alternative poly(A) signal sequences are present. The probable polyadenylation signals CATAAA or ACTAAA (HuPON1), AATACA (HuPON2), and AG-TAAA (HuPON3) are thought to be polyadenylated and cleaved inefficiently (Sheets et al., 1990). It may be relevant that Gieselmann et al. (1989) found that individuals with arylsulfatase A pseudodeficiency had a point mutation of the polyadenylation signal of the arylsulfatase A gene, which resulted in a substantial reduction in the amount of normal message. The amount of arylsulfatase protein and arylsulfatase enzyme activity was reduced 90% in individuals with arylsulfatase A pseudodeficiency. A second mutation affecting a glycosylation site was present in individuals with arylsulfatase A pseudodeficiency, but was found not to affect enzyme activity. Variation up to 13-fold in paraoxonase/arylesterase enzyme activity between individuals with the same allele type (e.g., homozygous low paraoxonase activity) has been observed (Furlong et al., 1989), and the levels observed are stable over time. It remains to be determined whether variations in polyadenylation signals between individuals of a given allele type contribute to the observed stable differences in enzyme levels. Alterations in the 5' regulatory region and stable differences in transcription factor levels could also

contribute to or be responsible for these differences.

The physiological substrate for paraoxonase has not been identified.

ACKNOWLEDGMENTS

We thank Drs. A. G. Motulsky and G. S. Omenn for encouragement and stimulating discussions throughout this project. Oligonucleotide synthesis was carried out by Susan Ribiero in the Chemistry Department or by the University of Washington Molecular Pharmacology Facility.

REFERENCES

- Aruffo, A., & Seed, B. (1987) *EMBO J.* 6, 3313-3316.
- Azuma, C., Tanabe, T., Konishi, M., Kinashi, T., Noma, T., Matsuda, F., Yaoita, Y., Takatsu, K., Hammarstrom, L., Smith, C. I. E., Severinson, E., & Honjo, T. (1986) *Nucleic Acids Res.* 14, 9149-9158.
- Blin, N., & Stafford, D. W. (1976) *Nucleic Acids Res.* 3, 2303-2308.
- Bock, S. C., Skriver, K., Nielsen, E., Thøgersen, H. C., Wiman, B., Donaldson, V. H., Eddy, R. L., Marrinan, J., Radziejewska, E., Huber, R., Shows, T. B., & Magnusson, S. (1986) *Biochemistry* 25, 4292-4301.
- Brown, S. S., Kalow, W., Pilz, W., Whittaker, M., & Waronick, C. L. (1981) *Adv. Clin. Chem.* 22, 1-123.
- Costa, L. G., Richter, R. J., Murphy, S. D., Omenn, G. S., Motulsky, A. G., & Furlong, C. E. (1987) in *Toxicology of Pesticides: Experimental, Clinical and Regulatory Perspectives* (Costa, L. G., Gall, C. L., & Murphy, S. D., Eds.), pp 262-266, Springer-Verlag, Heidelberg.
- Costa, L. G., McDonald, B. E., Murphy, S. D., Omenn, G. S., Richter, R. J., Motulsky, A. G., & Furlong, C. E. (1990) *Toxicol. Appl. Pharmacol.* 103, 66-76.
- Crabb, J. W., Armes, L. G., Carr, S. A., Johnson, C. M., Roberts, G. D., Bordoli, R. S., & McKeehan, W. L. (1986) *Biochemistry* 25, 4988-4993.
- Crabb, J. W., Johnson, C. M., Carr, S. A., Armes, L. G., & Saari, J. C. (1988) *J. Biol. Chem.* 263, 18678-18687.

- Furlong, C. E., Richter, R. J., Seidel, S. L., & Motulsky, A. G. (1988) *Am. J. Hum. Genet.* **43**, 230-238.
- Furlong, C. E., Richter, R. J., Seidel, S. L., Costa, L. G., & Motulsky, A. G. (1989) *Anal. Biochem.* **180**, 242-247.
- Furlong, C. E., Richter, R. J., Chapline, C., & Crabb, J. W. (1991) *Biochemistry* (preceding paper in this issue).
- Gan, K. N., Smolen, A., Eckerson, H. W., & LaDu, B. N. (1991) *Drug Metab. Dispos.* **19**, 100-106.
- Geldmacher-von Mallinckrodt, M., & Diepgen, T. L. (1988) *Toxicol. Environ. Chem.* **18**, 79-196.
- Gieselmann, V., Polten, A., Kreysing, J., & von Figura, K. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**, 9436-9440.
- Ginsburg, D., Zeheb, R., Yang, A. Y., Rafferty, U. M., Andreason, P. A., Nielsen, L., Dano, K., Lebo, R. V., & Gelehrter, T. D. (1986) *J. Clin. Invest.* **78**, 1673-1680.
- Harnik-Ort, V., Prakash, K., Marcantonio, E., Colman, D. R., Rosenfeld, M. G., Adesnik, M., Sabatini, D. D., & Kreibich, G. (1987) *J. Cell. Biol.* **104**, 855-863.
- Hassett, C., & Omiecinski, C. J. (1987) *Biochem. Biophys. Res. Commun.* **149**, 326-333.
- Hassett, C., & Omiecinski, C. J. (1990) *Nucleic Acids Res.* **18**, 1429-1434.
- Hassett, C., Turnblom, S. M., DeAngelis, A., & Omiecinski, C. J. (1989) *Arch. Biochem. Biophys.* **271**, 380-389.
- Hopp, T. P., & Woods, K. R. (1981) *Proc. Natl. Acad. Sci. U.S.A.* **78**, 3824-3828.
- Kalow, W. (1987) *Eur. J. Pharmacol.* **31**, 633-641.
- Knox, C. A. P., Sonthayanon, B., Chandra, G. R., & Muthukrishnan, S. (1987) *Plant Mol. Biol.* **9**, 3-17.
- Kvist, S., Roberts, L., & Dobberstein, B. (1983) *EMBO J.* **2**, 245-254.
- LaDu, B. N., & Eckerson, H. W. (1984) in *Genetic Variability in Response to Chemical Exposures* (Omenn, G. S., & Gelboin, H. V., Eds.) pp 167-175, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- LaDu, B. N., Piko, J. I., Eckerson, H. W., Vincent-Viry, M., & Seist, G. (1986) *Ann. Biol. Clin.* **44**, 369-372.
- Lalanne, J. L., Delarbre, C., Gachelin, G., & Kourilsky, P. (1983) *Nucleic Acids Res.* **11**, 1567-1577.
- Lycett, G. W., Croy, R. R. D., Shirsat, A. H., & Boulter, D. (1984) *Nucleic Acids Res.* **12**, 4493-4506.
- Main, A. R. (1956) *Can. J. Biochem. Physiol.* **34**, 197-216.
- Malissen, M., Malissen, B., & Jordan, B. R. (1982) *Proc. Natl. Acad. Sci. U.S.A.* **79**, 893-897.
- Meyer, W. E., Jonker, M., Klein, D., Ivanyi, P., van Seventer, G., & Klein, J. (1988) *EMBO J.* **7**, 2765-2774.
- Omenn, G. S. (1987) in *Toxicology of Pesticides: Experimental, Clinical and Regulatory Perspectives* (Costa, L. G., Galli, C. L., & Murphy, S. D., Eds.) pp 262-266, Springer-Verlag, Heidelberg.
- Omiecinski, C. J., Walz, F. G., Jr., & Vlasuk, G. P. (1985) *J. Biol. Chem.* **260**, 3247-3250.
- Pannekoek, H., Veerman, H., Lambers, H., Diergaarde, P., Verweij, C. L., van Zonneveld, A. J., & van Mourik, J. A. (1986) *EMBO J.* **5**, 2539-2544.
- Raines, C. A., Lloyd, J. C., Longstaff, M., Bradley, D., & Dyer, T. (1988) *Nucleic Acids Res.* **16**, 7931-7942.
- Sanger, F., Nicklen, S., & Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. U.S.A.* **74**, 5463-5467.
- Seidegard, J., Vorachek, W. R., Pero, R. W., & Pearson, W. R. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 7293-7297.
- Sheets, M. D., Ogg, S. C., & Wickens, M. P. (1990) *Nucleic Acids Res.* **18**, 5799-5805.
- Smolen, A., Eckerson, H. W., Gan, K. N., Hallat, N., & LaDu, B. N. (1991) *Drug Metab. Dispos.* **19**, 107-112.
- Sodoyer, R., Damotte, M., Delovitch, T. L., Trucy, J., Jordan, B. R., & Strachan, T. (1984) *EMBO J.* **3**, 879-885.
- Stevens, R. L., Avraham, S., Gartner, M. C., Bruns, G. A. P., Austen, K. F., & Weis, J. H. (1988) *J. Biol. Chem.* **263**, 7287-7291.
- Strandberg, L., Lawrence, D., & Ny, T. (1988) *Eur. J. Biochem.* **176**, 609-616.
- Takeda, J., Seino, Y., Tanaka, K. I., Fukumoto, H., Kayano, T., Takahashi, H., Mitani, T., Kurono, M., Suzuki, T., Tobo, T., & Imura, H. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 7005-7008.
- Tykocinski, M. L., Marche, P. N., Max, E. E., & Kindt, T. J. (1984) *J. Immunol.* **133**, 2261-2269.
- von Heijne, G. (1983) *Eur. J. Biochem.* **133**, 17-21.
- von Heijne, G. (1986) *Nucleic Acids Res.* **14**, 4682-4690.
- Watson, M. E. E. (1984) *Nucleic Acids Res.* **12**, 5145-5164.
- Weber, W. W. (1987) *The Acetyltransferase Genes and Drug Response*, Oxford University Press, New York.
- West, K., & Crabb, J. W. (1990) in *Current Research in Protein Chemistry* (Villafranca, J. J., Ed.) pp 37-48, Academic Press, San Diego.

GenBank Entry for PON1

LOCUS NM_000446 2395 bp mRNA linear PRI 21-JAN-2007

DEFINITION Homo sapiens paraoxonase 1 (PON1), mRNA.

ACCESSION NM_000446

VERSION NM_000446.3 GI:31880793

KEYWORDS .

SOURCE Homo sapiens (human)

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 2395)

AUTHORS Mastroianni, M., Mackness, M. and Mackness, B.

TITLE Defective metabolism of oxidized phospholipid by HDL from people with type 2 diabetes

JOURNAL Diabetes 55 (11), 3099-3103 (2006)

PUBMED 17065348

REMARK GeneRIF: In the control group, there was a significant negative correlation between serum PON1 activity and oxidized LDL concentration, but not in people with type 2 diabetes.

REFERENCE 2 (bases 1 to 2395)

AUTHORS Gaidukov, L., Rosenblat, M., Aviram, M. and Tawfik, D.S.

TITLE The 192R/Q polymorphs of serum paraoxonase PON1 differ in HDL binding, lipolactonase stimulation, and cholesterol efflux

JOURNAL J. Lipid Res. 47 (11), 2492-2502 (2006)

PUBMED 16914770

REMARK GeneRIF: PON1 192R/Q polymorphs differ in HDL binding, lipolactonase stimulation, and cholesterol efflux

REFERENCE 3 (bases 1 to 2395)

AUTHORS Hofer, S.E., Bennetts, B., Chan, A.K., Holloway, B., Karschinkus, C., Jenkins, A.J., Silink, M. and Donaghy, K.C.

TITLE Association between PON 1 polymorphisms, PON activity and diabetes complications

JOURNAL J. Diabetes Complicat. 20 (5), 322-328 (2006)

PUBMED 16949520

REMARK GeneRIF: Genotyping of adolescents with diabetes for PON 1 polymorphisms was performed, including that of a novel PON 1 promoter polymorphism A(-1074)G. PON genotypes were related to diabetes complication status.

REFERENCE 4 (bases 1 to 2395)

AUTHORS Lakshman, M.R., Gottipati, C.S., Narasimhan, S.J., Munoz, J., Marmillot, P. and Nylen, E.S.

TITLE Inverse correlation of serum paraoxonase and homocysteine thiolactonase activities and antioxidant capacity of high-density lipoprotein with the severity of cardiovascular disease in persons with type 2 diabetes mellitus

JOURNAL Metab. Clin. Exp. 55 (9), 1201-1206 (2006)

PUBMED 16919539

REMARK GeneRIF: cross-sectional study to correlate PON-1, homocysteine thiolactonase activities, and the lag time of LDL oxidation in control subjects and subjects with type 2 diabetes mellitus with different degrees of CVD

REFERENCE 5 (bases 1 to 2395)

AUTHORS Yamane, T., Matsumoto, T., Nakae, I., Takashima, H., Tarutani, Y., Tamaki, S. and Horie, M.

TITLE Impact of paraoxonase polymorphism (Q192R) on endothelial function in intact coronary circulation

JOURNAL Hypertens. Res. 29 (6), 417-422 (2006)

PUBMED 16940704

REMARK GeneRIF: as estimated by bradykinin and acetylcholine testing, findings suggest that paraoxonase-1 (PON1) genotypes may not play a critical role in the modulation of endothelial vasomotor function in the intact coronary circulation.

REFERENCE 6 (bases 1 to 2395)

AUTHORS Adkins, S., Gan, K.N., Mody, M. and La Du, B.N.

TITLE Molecular basis for the polymorphic forms of human serum paraoxonase/arylesterase: glutamine or arginine at position 191, for the respective A or B allozymes

JOURNAL Am. J. Hum. Genet. 52 (3), 598-608 (1993)

PUBMED 7916578
 REFERENCE 7 (bases 1 to 2395)
 AUTHORS Humbert,R., Adler,D.A., Distech,C.M., Hassett,C., Omiecinski,C.J. and Furlong,C.E.
 TITLE The molecular basis of the human serum paraoxonase activity polymorphism
 JOURNAL Nat. Genet. 3 (1), 73-76 (1993)
 PUBMED 8098250
 REFERENCE 8 (bases 1 to 2395)
 AUTHORS Hassett,C., Richter,R.J., Humbert,R., Chapline,C., Crabb,J.W., Omiecinski,C.J. and Furlong,C.E.
 TITLE Characterization of cDNA clones encoding rabbit and human serum paraoxonase: the mature protein retains its signal sequence
 JOURNAL Biochemistry 30 (42), 10141-10149 (1991)
 PUBMED 1657140
 REFERENCE 9 (bases 1 to 2395)
 AUTHORS Gan,K.M., Smolen,A., Eckerson,H.W. and La Du,B.N.
 TITLE Purification of human serum paraoxonase/arylesterase. Evidence for one esterase catalyzing both activities
 JOURNAL Drug Metab. Dispos. 19 (1), 100-106 (1991)
 PUBMED 1673382
 REFERENCE 10 (bases 1 to 2395)
 AUTHORS Ortigoza-Ferado,J., Richter,R.J., Hornung,S.K., Motulsky,A.G. and Furlong,C.E.
 TITLE Paraoxon hydrolysis in human serum mediated by a genetically variable arylesterase and albumin
 JOURNAL Am. J. Hum. Genet. 36 (2), 295-305 (1984)
 PUBMED 6324579
 COMMENT VALIDATED REFSEQ: This record has undergone preliminary review of the sequence, but has not yet been subject to final review. The reference sequence was derived from [CB122285.1](#) and [Z70723.1](#). On Jun 18, 2003 this sequence version replaced [gi:19923105](#).

Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

FEATURES
 source 1..2395
 /organism="Homo sapiens"
 /mol_type="mRNA"
 /db_xref="taxon:9606"
 /chromosome="7"
 /map="7q21.3"
 gene 1..2395
 /gene="PON1"
 /note="paraoxonase 1; synonyms: ESA, PON"
 /db_xref="GeneID:5444"
 /db_xref="HGNC:9204"
 /db_xref="HPRD:01351"
 /db_xref="MIM:168820"
 STS 29..1156
 /gene="PON1"
 /db_xref="UniSTS:481927"
 CDS 58..1125
 /gene="PON1"
 /EC_number="3.1.1.2"
 /EC_number="3.1.8.1"
 /go_component="extracellular region [pmid 14718574]"
 /go_function="aryldialkylphosphatase activity;
 arylesterase activity [pmid 1673382]; hydrolase activity;
 lipid transporter activity"
 /go_process="response to external stimulus [pmid 1657140]"
 /note="Paraoxonase; esterase A"
 /codon_start=1
 /product="paraoxonase 1"
 /protein_id="NP_000437.3"
 /db_xref="GI:19923106"
 /db_xref="CCDS:CCDS5638.1"
 /db_xref="GeneID:5444"
 /db_xref="HGNC:9204"
 /db_xref="HPRD:01351"

compare to
sequence in
Fig.1A and 1B.
Doesn't match.

```

/db_xref="MIM:168820"
/translation="MKKLIALTLLMGLALPRNHQSSYQTRLNALREVQPVELPNCNL
VKGIFTGSEDLLEILPGLAFISSGLKYPGLKSPNPNPSGKILLMDLNEEDPTVLELGI
TGSKPDVSSFPNHGISTPTDENDMAYLVIVNHPDAKSTVELFKPQREKESLLHLKTIK
HKLILPNLNDIVAVGPRHPYGTNDHYFLDPYLQSWENYGLAWSVYVYSPSEVRVVAE
GFDPFANGINISPDGKYVYIAELLAHKIHVYERHANWTLTPLKSLDFNTLVNDSVDPE
TGDLMVGCHPNGMKIIFYDSENPFASEVLRIQNILTEEPKVTQVYAENGTVLQGSTVA
SVYKGLLIGTVPHKALYCEL"
557..670
/gene="PONI"
/standard_name="G44327"
/db_xref="UniSTS:95116"
564..662
/gene="PONI"
/standard_name="GDB:194296"
/db_xref="UniSTS:155789"
1132..1269
/gene="PONI"
/standard_name="STS-M63012"
/db_xref="UniSTS:46466"
1441..1619
/gene="PONI"
/standard_name="RH12859"
/db_xref="UniSTS:63408"

ORIGIN
1 agagcctcct agccctctgg tgtctgcgcc catcgatccc ttgtctctac cccgaccattg
61 gcgaagctga ttgcgctcac cctcttgggg atgggactgg cactcttcag gaaccaccagg
121 tctctctacc aaacacgact taatgctctc cgagaggtac aaccctaga actctctaac
181 tgaattattg ttaaggaat cgaaactggc tctgaagact tggagatact gcttaattgga
241 ctggctttca ttatgctctgg attaaagtat cctggaaata agagcttcaa ccccaacagt
301 cctggaaaaa tacttctgat ggacctgaat gaagaagatc caacagtgtt ggaattgggg
361 atcactggaa gtaaaattga tgtatcttca tttaaacctc atgggattag cacatctaca
421 gatgaagata atgcactgta cctcctgggt gtgaacctac cagatgccaa gtccacagtg
481 gagttgttta aatttcaga agaagaaaaa tctgttttgc atctaaaaac catcagacat
541 aaactctctg ctaatttgaa tgaattgttt gctgtgggag ctgagaoact ttatggaca
601 aatgcactac attttcttga cccctactta caactctggg agatgtattt gggtttagcg
661 tggctgtatg ttgtctacta tagtccaagt gaagtctgag tgggtggcga aggattgat
721 ttgtctaatg gaatcaacat ttcaaccgat ggcaagtatg tctatatagc tgagtttgtg
781 gctcataaga ttcatgtgta tgaaaagcat gctaatttga ctttaactcc attgaagtcc
841 cttgacttta ataccctctg ggataacata ttctgtgata ctgacagagg agacctttgg
901 gttggatgcc atcccaatgg catgaaaatc ttctctatg actcaagaaa tctccttgca
961 tcagaggtgc ttogaatcca gaacattcta acagaagaac ctaaagtgac acaggtttat
1021 gcagaaaaatg gcacagtggt gcaaggcagt acagttgcct ctgtgtacaa agggaaaactg
1081 ctgatttgga cagtgtttca caaagctctt tactgtgagc tctaacagac cgatttgcac
1141 ccatgccata gaaactgagg ccattatttc aaccgcttgc catattccga ggacccagtg
1201 tctttagctg aacaatgaat gctgacctta aatgtggaca tcatggaaga tcaaaagcact
1261 gtttaactgg gagtgatag atgtgtgggg ctttttttgg agaatacaat atcaaatcaag
1321 tcttgaataa cttgaaaaac ctaattacca taaaaatcct tctactaaaa atggataaat
1381 cagttatgtc aattgtcaga tattaaataa caagtgtgga ccccaaaagt acttaacctc
1441 aaacatgtgt tgcctgaaag cacatgtgtg tactgctgoc ttgcatgic ttgttcagaa
1501 gacacaggggg agcaggggta gctcaagtgct ctttagaact ccaagtactca cccagggact
1561 ccagttcaaca ggcacagaaaa catatgaatt atgaagttcc cctctactcc atgacatag
1621 taagtctgac tatggcagtc agacttactt actccacttt tcoctctgat atatgacttt
1681 tctcagtaaa atattaaact gaactattcc aactccoctt gtactcttgc tttttcaatt
1741 ctctctgtgc aatgacacat agggaaaatc taaaattctt gggagtggtg tcaacactga
1801 attgattgag tctctatgat cttggcacaa attgtacatt tgaagtcttt tgacttgggt
1861 aaagggaagt ttgtcaacttc gatgactgga tacagaatga atcccaaat tgacatgggg
1921 gcagtgaaaa gtgtccoccaa agactacaact gttgtgagg tgggtggtagt gacttgggt
1981 ttttgtttaa tatttaaaact tctgtgttgc gagggtgaaa agaaaaaaaaa ggcatactac
2041 gtaaacaaac aaataaatag aaaaagatcaa caacccctttt ggctatctac tgagacatga
2101 ctaggagaag aaatgacttt tatcattttg ttatagaagg tgatatataa ggttacacat
2161 ttctcattat ttgttttttt gatttgaagg tataaccttc atgatgaatt actctctcag
2221 ggtgttaagg cagtgaacttt agaacaacaa tttttctttt tttttttttt gtttttgaga
2281 ccgaattcca ctctgttgcc caggctggag tgcagttgtg cgtactttgc tcaactgcaac
2341 ttctacotcc gaggttcaag agattcttgt gctccagoct cccgtagatg tgcg

```

//

GenBank entry for Pon2 (the coding sequence in Patent US 5,629,193)

LOCUS NM_000305 1669 bp mRNA linear PRI 03-DEC-2006
 DEFINITION Homo sapiens paraoxonase 2 (PON2), transcript variant 1, mRNA.
 ACCESSION NM_000305
 VERSION NM_000305.2 GI:66529293
 KEYWORDS .
 SOURCE Homo sapiens (human)
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini;
 Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 1669)
 AUTHORS Ng, C.J., Bourquard, N., Grijalva, V., Hama, S., Shih, D.M., Navab, M.,
 Fogelman, A.M., Lusis, A.J., Young, S. and Reddy, S.T.
 TITLE Paraoxonase-2 deficiency aggravates atherosclerosis in mice despite
 lower apolipoprotein-B-containing lipoproteins: anti-atherogenic
 role for paraoxonase-2
 J. Biol. Chem. 281 (40), 29491-29500 (2006)
 PUBMED 16891303
 REMARK GeneRIF: Plays a protective role against atherosclerosis in vivo.
 REFERENCE 2 (bases 1 to 1669)
 AUTHORS Slowik, A., Tomik, B., Wolkow, P.P., Partyka, D., Turaj, W.,
 Malecki, M.T., Pera, J., Dziedzic, T., Szczudlik, A. and Figlewicz, D.A.
 TITLE Paraoxonase gene polymorphisms and sporadic ALS
 J. Neurology 67 (5), 766-770 (2006)
 PUBMED 16822965
 REMARK GeneRIF: the C311S polymorphism was associated with sALS in
 dominant and additive models.
 REFERENCE 3 (bases 1 to 1669)
 AUTHORS Saeed, M., Siddique, N., Rung, W.Y., Usacheva, E., Liu, E., Sufit, R.L.,
 Heller, S.L., Haines, J.L., Pericak-Vance, M. and Siddique, T.
 TITLE Paraoxonase cluster polymorphisms are associated with sporadic ALS
 J. Neurology 67 (5), 771-776 (2006)
 PUBMED 16522964
 REMARK GeneRIF: A haplotype of high linkage disequilibrium (LD) spanning
 PON2 and PON3 was associated with SALS.
 REFERENCE 4 (bases 1 to 1669)
 AUTHORS Dragancov, D.I., Teiber, J.F., Speelman, A., Osawa, Y., Sunahara, R. and
 La Du, B.N.
 TITLE Human paraoxonases (PON1, PON2, and PON3) are lactonases with
 overlapping and distinct substrate specificities
 J. Lipid Res. 46 (6), 1239-1247 (2005)
 JOURNAL 15772423
 PUBMED 15772423
 REMARK GeneRIF: PON1, PON2, and PON3 are lactonases with overlapping and
 distinct substrate specificities
 REFERENCE 5 (bases 1 to 1669)
 AUTHORS Ng, C.J., Shih, D.M., Hama, S.Y., Villa, N., Navab, M. and Reddy, S.T.
 TITLE The paraoxonase gene family and atherosclerosis
 Free Radic. Biol. Med. 38 (2), 153-163 (2005)
 JOURNAL 15607899
 PUBMED 15607899
 REMARK Review article
 REFERENCE 6 (bases 1 to 1669)
 AUTHORS Ng, C.J., Wadleigh, D.J., Gangopadhyay, A., Hama, S., Grijalva, V.R.,
 Navab, M., Fogelman, A.M. and Reddy, S.T.
 TITLE Paraoxonase-2 is a ubiquitously expressed protein with antioxidant
 properties and is capable of preventing cell-mediated oxidative
 modification of low density lipoprotein
 J. Biol. Chem. 276 (48), 44444-44449 (2001)
 JOURNAL 11579088
 PUBMED 11579088
 REFERENCE 7 (bases 1 to 1669)
 AUTHORS Hong, S.H., Song, J., Min, W.K. and Kim, J.Q.
 TITLE Genetic variations of the paraoxonase gene in patients with
 coronary artery disease
 Clin. Biochem. 34 (6), 475-481 (2001)
 JOURNAL 11676977
 PUBMED 11676977
 REMARK GeneRIF: study suggested a gene-gene interaction between the PON1
 and PON2 polymorphisms for CAD risk; may have linkage
 disequilibrium with a tightly linked PON3 locus or significant
 atherosclerotic alleles of nearby genes

REFERENCE 8 (bases 1 to 1669)
AUTHORS Mochizuki,H., Scherer,S.W., Xi,T., Nickle,D.C., Majer,M.,
Ruisenga,J.J., Tsui,L.C. and Prochazka,M.
TITLE Human PON2 gene at 7q21.3: cloning, multiple mRNA forms, and
missense polymorphisms in the coding sequence
JOURNAL Gene 213 (1-2), 149-157 (1998)
PUBMED 9714608

REFERENCE 9 (bases 1 to 1669)
AUTHORS Sanghera,D.K., Aston,C.E., Saha,N. and Kamboh,M.I.
TITLE DNA polymorphisms in two paraoxonase genes (PON1 and PON2) are
associated with the risk of coronary heart disease
JOURNAL Am. J. Hum. Genet. 62 (1), 36-44 (1998)
PUBMED 9443862

REFERENCE 10 (bases 1 to 1669)
AUTHORS Primo-Farmo,S.L., Sorenson,R.C., Teiber,J. and La Du,B.N.
TITLE The human serum paraoxonase/arylesterase gene (PON1) is one member
of a multigene family
JOURNAL Genomics 33 (3), 498-507 (1996)
PUBMED 8661009

COMMENT REVIEWED REFSEQ: This record has been curated by NCBI staff. The
reference sequence was derived from [CB961097.1](#), [AF001601.1](#) and
[BC040010.1](#).
On May 24, 2005 this sequence version replaced [gi:4505952](#).

Summary: This gene encodes a member of the paraoxonase gene family,
which includes three known members located adjacent to each other
on the long arm of chromosome 7. The encoded protein is
ubiquitously expressed in human tissues, membrane-bound, and may
act as a cellular antioxidant, protecting cells from oxidative
stress. Hydrolytic activity against acylhomoserine lactones,
important bacterial quorum-sensing mediators, suggests the encoded
protein may also play a role in defense responses to pathogenic
bacteria. Mutations in this gene may be associated with vascular
disease and a number of quantitative phenotypes related to
diabetes. Alternatively spliced transcript variants encoding
different isoforms have been described.

Transcript Variant: This variant (1) represents the longer
transcript and encodes the longer isoform (1).

Publication Note: This RefSeq record includes a subset of the
publications that are available for this gene. Please see the
Entrez Gene record to access additional publications.
COMPLETENESS: complete on the 3' end.

FEATURES
source Location/Qualifiers
1..1669
/organism="Homo sapiens"
/mol_type="mRNA"
/db_xref="taxon:9606"
/chromosome="7"
/map="7q21.3"

gene 1..1669
/gene="PON2"
/note="paraoxonase 2"
/db_xref="GeneID:5445"
/db_xref="HGNC:9205"
/db_xref="HPRD:03903"
/db_xref="MIM:602447"

CDS 122..1186
/gene="PON2"
/EC_number="3.1.1.2"
/EC_number="3.1.8.1"
/go_component="extracellular region; membrane"
/go_function="aryldialkylphosphatase activity;
arylesterase activity [pmid 8661009]; hydrolase activity"
/note="isoform 1 is encoded by transcript variant 1; serum
paraoxonase/arylesterase 2; aromatic esterase 2;
A-esterase 2; serum aryldialkylphosphatase 2; paraoxonase
nirs"
/codon_start=1
/product="paraoxonase 2 isoform 1"

compare to
sequence in
Fig.1A and 1B.
Perfect match.

```

/protein id="NP_000296.2"
/db_xref="GI:66529294"
/db_xref="CCDS:CCDS5640.1"
/db_xref="GeneID:5445"
/db_xref="HGNC:9205"
/db_xref="HPRD:03903"
/db_xref="MIM:602447"
/translation="MGRILAVGLLGIALALLGERILLARNLKASREVESVDLPCHL
IKIGRAGSEDIIDILPNGLAFFSVGLKFPGLHSFAPDKPGIIMMDLKEEKPRARELRI
SRGFDLASFPNPHIGI STFIIDNDITVYLFVNVHPEFKNTVEIKFPEEAENSLHLKTVKH
ELLPSVNDITAVGPAHFYATNDHYFSDPFLKYLETYLNLHWANVVYSPNEVKVVAEG
FDSANGINISPDKKIYVADILAHEHIVLEKHTNMNLTQLKVELEDTLVNLSIDPSS
GDIHVGCHPNGKLFVYDPNPPSFSEVLRIRQNLISEKPTVTITVYANNQSVLQGSVAS
VYDGKLLIGTLYHRYALYCEL"
1145..1278
/gene="PON2"
/standard_name="D7S2922"
/db_xref="UniSTS:73923"
1385..1560
/gene="PON2"
/standard_name="D7S2091E"
/db_xref="UniSTS:2676"
1439..1558
/gene="PON2"
/standard_name="D7S2405"
/db_xref="UniSTS:154285"
1632..1637
/gene="PON2"
1654
/gene="PON2"

```

STS

STS

STS

polyA signal

polyA site

ORIGIN

```

1   ggactcggcc   taggcggagg   acggggcgga   ggcggccggg   caccatcgag   cggggaagat
61  ggcaccgccc   acggagctgc   tggccaggcc   ggagcgagcc   agcgcgcccg   gctccgcgcg
121 catggggcgg   ctgggtgcgt   tgggcttgct   gggggtcgcg   ctggcgctcc   tgggcgagag
181 gcttctggca   ctcaaaaatc   gaactaaagc   ctccagagaa   gtagaatctg   tagacctccc
241 acactgccac   ctgattaaag   gaattgaagc   tggctctgaa   gatattgaca   tacttcccaa
301 tggctcggct   ttttttagtg   tgggtctaaa   attcccagga   ctccacagct   ttgcaccaga
361 taagcctcga   ggaatactaa   tgaaggatct   aaagaagaa   aaaccaaggc   ccacgggaat
421 aagatcagct   cgtgggttga   atttggcttc   attcaatcca   catggcatca   gcaattcact
481 agacaacgat   gacacagttt   atctctttgt   tgtaaaacc   ccagatctca   agaatacagt
541 gcaaatTTTT   aaatttgaag   aagcagaaaa   ttctctgttg   catctgaaaa   cagtcaaaica
601 tgaagctctt   ccaagtgtga   atgacatcac   agctgttgga   ccggcacatt   tctatgccac
661 aaatgacacc   tacttctctg   atcctttctt   aaagtattta   gaaacatact   tgaacttaca
721 ctggcgaaat   gttgtttact   acagtccaaa   tgaagttaaa   gtggtagcag   aaggatttga
781 ttcagcaaat   gggatcaata   ttccacctga   tgataagtat   atctatgttg   ctgacatat
841 ggctcatgaa   attcatgttt   tggaaaaaca   cactaatatg   aatttaactc   agttgaaggt
901 acttgagctg   gatacactgg   ttgataattt   atctattgat   ccttctctgg   gggacatctg
961 ggtgagcgtg   catcctaagt   gccagaagct   ctctgtgtat   gacccgaaac   atcctccctc
1021 gtccagaggtt   ctccgcctcc   agaacattct   atctgagaag   cctacagtga   ctacagttta
1081 tgcaccaaat   gggctctgtc   tccaaggag   ttctgtagcc   tcagtgtagt   atggggaagt
1141 gctcatagcg   accttatacc   acagagcctt   gtattgtgaa   ctcaaatgtg   tacttttggc
1201 atgaaagtgc   gataacctaa   caattaattt   tctatgaatt   gctaattctg   agggaaattt
1261 accagcaaca   tgcacocaga   aactgatggc   atcggtagtt   aatttattc   cagttaagaa
1321 cggccctttt   agtcctttag   gtccttttaa   caaaaaagga   aatttattc   cagttaagaa
1381 atgccaaaga   agggacagaa   aagaagctgc   ctttcgata   aagtgaatac   attttgaca
1441 aagtaagcct   caactttgoc   ttccaaactg   cagaacatgg   attccactga   aatagagtga
1501 attatatttc   cttaaaatgt   gagtgaactc   acttctgcca   ctgtgactac   tatggctgtt
1561 tagaactact   gataacgtat   ttgatgttt   tgtacttaca   tctttgttta   coattaaaaa
1621 gctggagtta   tattaaagac   taactaaaat   cccaaaaaaa   aaaaaaaa

```